arXiv:1012.4971v1 [quant-ph] 22 Dec 2010

Patterns in Wigner-Weyl approach

Antonina N. Fedorova, Michael G. Zeitlin

Mathematical Methods in Mechanics Group
Institute of Problems of Mechanical Engineering (IPME RAS)
Russian Academy of Sciences
Russia, 199178, St. Petersburg, V.O., Bolshoj pr., 61
zeitlin@math.ipme.ru, anton@math.ipme.ru
http://www.ipme.ru/zeitlin.html
http://mp.ipme.ru/zeitlin.html

Abstract

We present a family of methods, which can describe behaviour of quantum ensembles and demonstrate the creation of nontrivial (meta) stable states (patterns), localized, chaotic, entangled or decoherent from basic localized modes in collective models arising from the quantum hierarchy of Wigner-von Neumann-Moyal equations.

Presented: GAMM Meeting, 2006, Berlin, Germany.

Published: Proc. Appl. Math. Mech. (PAMM), **6**, 627-628, Wiley-VCH, 2006.

DOI 10.1002/pamm.200610294

Patterns in Wigner-Weyl approach

Antonina N. Fedorova and **Michael G. Zeitlin**[*]

IPME RAS, St. Petersburg, V.O. Bolshoj pr., 61, 199178, Russia

We present a family of methods, which can describe behaviour of quantum ensembles and demonstrate the creation of nontrivial (meta) stable states (patterns), localized, chaotic, entangled or decoherent from basic localized modes in collective models arising from the quantum hierarchy of Wigner-von Neumann-Moyal equations.

Copyright line will be provided by the publisher

It is widely known that the currently available experimental techniques in the area of quantum physics as a whole and in that of quantum computations in particular, as well as the present level of understanding of phenomenological models, outstripped the actual level of mathematical/theoretical description. Considering, for example, the problem of describing the realizable states one should not expect that planar waves and (squeezed) gaussian coherent states would be enough to characterize such complex systems as qCPU (quantum Central Processor Unit)-like devices. Complexity of the set of relevant states, including entangled (chaotic) states is still far from being clearly understood and moreover from being realizable. Our main motivations arise from the following general questions: How can we represent well localized and reasonable state in mathematically correct form? Is it possible to create entangled and other relevant states by means of these new building blocks? A starting point for us is a possible model for (continuous) "qudit" with subsequent description of the whole zoo of possible realizable (controllable) states/patterns which may be useful from the point of view of quantum experimentalists and engineers. Effects we are interested in are: (a)hierarchy of internal/hidden scales (time, space, phase space); (b)non-perturbative multiscales: from slow to fast contributions, from the coarser to the finer level of resolution/decomposition; (c)coexistence of hierarchy of multiscale dynamics with transitions between scales. (d)realization of the key features of the complex quantum world such as the existence of chaotic and/or entangled states with possible destruction in "open/dissipative" regimes due to interactions with quantum/classical environment and transition to decoherent states. By localized states (localized modes) we mean the building blocks for solutions or generating modes which are localized in maximally small region of the phase (as in c-(classical) as in q(quantum)-case) space. By an entangled/chaotic pattern we mean some solution (or asymptotics of solution) which has random-like distributed energy (or information) spectrum in a full domain of definition. In quantum case we need to consider additional entangled-like patterns, roughly speaking, which cannot be separated into pieces of sub-systems. By a localized pattern (waveleton) we mean (asymptotically) (meta) stable solution localized in a relatively small region of the whole phase space (or a domain of definition). In this case the energy is distributed during some time (sufficiently large) between only a few localized modes (from point 1). We believe it to be a good model for plasma in a fusion state (energy confinement) or a model for quantum continuous "qubit" or a result of the decoherence process in open quantum system when the full entangled state degenerates into localized (quasiclassical) pattern. In this paper we consider the calculations of the Wigner functions $W(p,q,t)$ (WF) corresponding to the classical polynomial Hamiltonian $H(p,q,t)$ as the solution of the Wigner-von Neumann equation: $i\hbar W_t = H * W - W * H$ and related Wigner-like equations for different ensembles. According to the Weyl transform, a quantum state (wave function or density operator ρ) corresponds to the Wigner function, which is the analogue in some sense of classical phase-space distribution [1]. Wigner equation is a result of the Weyl transform or "wignerization" of von Neumann equation for density matrix. The difference between classical and quantum case is concentrated in the structure of the set of operators included in the set-up and, surely, depends on the method of quantization. But, in the naive Wigner-Weyl approach for quantum case the symbols of operators play the same role as usual functions in classical case. To move from c- to q-case, let us start from the second quantized representation for an algebra of observables $A = (A_0, A_1, \dots, A_s, ...)$ in the standard form $A = A_0 + \int dx_1 \Psi^+(x_1) A_1 \Psi(x_1) + \cdots + (s!)^{-1} \int dx_1 \dots dx_s \Psi^+(x_1) \dots \Psi^+(x_s) A_s \Psi(x_s) \dots \Psi(x_1) + \dots$ N-particle Wigner function $W_s(x_1, \dots, x_s) = \int dk_1 \dots dk_s \exp\big(-i \sum_{i=1}^s k_i p_i\big) \mathrm{Tr} \rho \Psi^+\big(q_1 - \frac{1}{2}\hbar k_1\big) \dots, \Psi^+\big(q_s - \frac{1}{2}\hbar k_s\big) \Psi\big(q_s + \frac{1}{2}\hbar k_s\big) \dots \Psi\big(q_1 + \frac{1}{2}\hbar k_s\big)$ allow us to consider them as some quasiprobabilities and provide useful bridge between c- and q-cases: $< A >= \mathrm{Tr} \rho A = \sum_{s=0}^\infty (s!)^{-1} \int \prod_{i=1}^s d\mu_i A_s(x_1, \dots, x_s) W_s(x_1, \dots, x_s)$. The full description for quantum ensemble can be done by the whole hierarchy of functions (symbols): $W = \{W_s(x_1, \dots, x_s), s = 0, 1, 2 \dots\}$ So, we may consider the following q-hierarchy as the result of "wignerization" procedure for c-BBGKY one, where the partial Wigner functions are solutions of proper Wigner equations:

$$\frac{\partial W_n}{\partial t} = -\frac{p}{m}\frac{\partial W_n}{\partial q} + \sum_{\ell=0}^{\infty} \frac{(-1)^\ell (\hbar/2)^{2\ell}}{(2\ell+1)!} \frac{\partial^{2\ell+1} U_n(q)}{\partial q^{2\ell+1}} \frac{\partial^{2\ell+1} W_n}{\partial p^{2\ell+1}}. \tag{1}$$

---

[*] Corresponding author: e-mail: zeitlin@math.ipme.ru, http://www.ipme.ru/zeitlin.html, http://mp.ipme.ru/zeitlin.html

Copyright line will be provided by the publisher



Fig. 1 Localized pattern or waveleton Wigner function (left); entangled-like Wigner function (right).

We obtain our multiscale/multiresolution representations for solutions of Wigner-like equations via a variational-wavelet approach. We represent the solutions as decomposition into localized eigenmodes (regarding action of affine group, i.e. hidden symmetry of the underlying functional space of states) related to the hidden underlying set of scales: $W_n(t,q,p) = \bigoplus_{i=i_c}^{\infty} W_n^i(t,q,p)$, where value i_c corresponds to the coarsest level of resolution c or to the internal scale with the number c in the full multiresolution decomposition (MRA) of the underlying functional space (L^2, e.g.): $V_c \subset V_{c+1} \subset V_{c+2} \subset \dots$ We introduce the Fock-like space structure (in addition to the standard one, if we consider second-quantized case) on the whole space of internal hidden scales $H = \bigoplus_i \bigotimes_n H_i^n$ for the set of n-partial Wigner functions (states): $W^i = \{W_0^i, W_1^i(x_1;t), \dots, W_N^i(x_1,\dots,x_N;t), \dots\}$, where $W_p(x_1,\dots,x_p;t) \in H^p$, $H^0 = C$, $\quad H^p = L^2(R^{6p})$ (or any different proper functional space), with the natural Fock space like norm: $(W,W) = W_0^2 + \sum_i \int W_i^2(x_1,\dots,x_i;t) \prod_{\ell=1}^{i} \mu_\ell$. We consider $W = W(t)$, $W \in L^2(R)$, via multiresolution decomposition which naturally and efficiently introduces the infinite sequence of the underlying hidden scales [2]. We have the contribution to the final result from each scale of resolution from the whole infinite scale of spaces. The closed subspace D_j(j$\in$ $\mathbf{Z}$) corresponds to the level j of resolution, or to the scale j. Then we have the following decomposition: $\{W(t)\} = \bigoplus_{-\infty<j<\infty} D_j = \overline{V_c \bigoplus_{j=0}^{\infty} D_j}$, in case when V_c is the coarsest scale of resolution. As a result the solution of this hierarchy has the following multiscale or multiresolution decomposition via nonlinear high-localized eigenmodes

$$W(t,x_1,x_2,\dots) = \sum_{(i,j)\in Z^2} a_{ij} U^i \otimes V^j(t,x_1,\dots), \tag{2}$$

$$V^j(t) = V_N^{j,slow}(t) + \sum_{l\geq N} V_l^j(\omega_l t),\ \omega_l \sim 2^l, \quad U^i(x_s) = U_M^{i,slow}(x_s) + \sum_{m\geq M} U_m^i(k_m^s x_s),\ k_m^s \sim 2^m,$$

which corresponds to the full multiresolution expansion in all underlying time/space scales. It should be noted that such representations give the best possible localization properties in the corresponding (phase)space/time coordinates. Numerical calculations are based on compactly supported wavelets and wavelet packets and on evaluation of the accuracy on the level N of the corresponding cut-off of the full system regarding Fock-like norm above: $\|W^{N+1} - W^N\| \leq \varepsilon$.

The numerical simulation demonstrates the formation of different (stable) pattern or orbits generated by internal hidden symmetry from high-localized structures. Our (nonlinear) eigenmodes are more realistic for the modeling of nonlinear classical/quantum dynamical process than the corresponding linear gaussian-like coherent states. Here we mention only the best convergence properties of the expansions based on wavelet packets, which realize the minimal Shannon entropy property and the exponential control of convergence of expansions like (2) based on the norm above. Fig. 1 (left) corresponds to (possible) result of superselection (einselection) [1] after decoherence process started from entangled state (Fig. 1, right).

References

[1] W. P. Schleich, Quantum Optics in Phase Space, (Wiley, 2000); D. Sternheimer, Deformation quantization, math/9809056.

[2] A.N. Fedorova and M.G. Zeitlin, in: *Quantum Aspects of Beam Physics*, Ed. P. Chen (World Scientific, Singapore, 2002) p. 527, 539; physics/0101006; physics/0101007; in: *Quantum Aspects of Beam Physics*, Eds. Pisin Chen, K. Reil (World Scientific, 2004) p 22; quant-phys/0306197; *Nuclear Instruments and Methods in Physics Research Section A*, **534**, Issues 1-2, 309, 314, (2004); quant-ph/0406009; quant-ph/0406010.

[3] A.N. Fedorova and M.G. Zeitlin, Int. J. of Modern Physics B, Vol. 20, 1570 (2006).

Copyright line will be provided by the publisher